

RESTRICTED USE PESTICIDE

Due to Acute Oral, Dermal and Inhalation Toxicity and Risks to Aquatic Organisms and Wildlife.

For retail sale to, and use only by Certified Applicators or persons under their direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

Counter[®] 15G

systemic insecticide nematicide

LOCK[™]inLOAD[™]

CLOSED LOADING SYSTEM

FOR USE IN FIELD CORN, FIELD CORN GROWN FOR SEED, POPCORN, SWEET CORN, SUGAR BEETS, AND GRAIN SORGHUM

ACTIVE INGREDIENT:	By Weight
Terbufos: S-[[[(1,1-dimethylethyl)thio]methyl]O,0-diethyl phosphorodithioate.....	15.0%
INERT INGREDIENTS	85.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
DANGER! PELIGRO!



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
Organophosphate (a cholinesterase inhibitor)	
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to poisoning. Have the product label with you when calling a poison control center or doctor or going for treatment. Antidote: Atropine is an antidote	
EMERGENCY INFORMATION	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call a physician in all cases of suspected poisoning. If warning symptoms appear, keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.	
FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:	
Transportation: CHEMTREC.....	1-800-424-9300
Other: AMVAC.....	1-323-264-3910

EPA Reg. No. 5481-545
EPA Est. No. 5905-1A-01
Made in U.S.A.

Net Contents:
As Marked on Container



4100 E. Washington Boulevard
Los Angeles, CA 90023, U.S.A.
1-323-264-3910 • www.amvac-chemical.com

NOTE TO PHYSICIAN

Warning signs and symptoms include weakness, muscle twitching, headache, tightness in chest, respiratory depression, blurred vision, miosis, hypersecretion, nausea, vomiting, diarrhea, and abdominal cramps. If poisoning is probable, treat the patient immediately. Clear airway and provide oxygen before administering atropine. Give atropine intravenously (IV), intramuscularly or through an alternative route such as an endotracheal tube. Depending on severity of poisoning, use 2 to 4 milligrams IV in adults (or 0.05 mg/kg in children under 12 years) every 10-15 minutes until fully atropinized as shown primarily by control of pulmonary secretions. Proceed concurrently with decontamination using proper protective gear; for example, chemical-resistant gloves rather than cotton or leather ones. Use soap (preferably, tincture of green soap) and water, or a dilute hypochlorite solution, for decontaminating skin. Maintain atropinization by repeated doses based on recurrence of symptoms. Twenty to thirty milligrams, or more, may be required during the first 24 hours, and preservative-free atropine preparation should be used if large doses are required. Generally, inducing emesis is not advisable. Avoid opiates, parasympathomimetic agents (e.g. succinylcholine), theophylline, reserpine, and or phenothiazines. Suction oral secretions and emesis to avoid aspiration. Artificial respiration or oxygen administration may be necessary. Probable mucosal damage may contraindicate the use of gastric lavage. If not contraindicated, gastric emptying by lavage may be considered after ingestion of a potentially life-threatening amount if it can be performed soon after ingestion (within an hour). A slurry of activated charcoal may be medically warranted at an oral dose appropriate for adult, child, or infant. For seizures, give a benzodiazepine intravenously. Observe patient continuously for at least 72 hours. Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) appears to be an effective adjunct to atropine, especially in those severely ill and if given soon. Before it's administered, obtain a blood sample for cholinesterase analysis. Adjusting for age and weight, pralidoxime may be administered as a continuous infusion after a loading dose or using a bolus method.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER!

Fatal if swallowed, inhaled, or absorbed through the skin. Corrosive. Causes irreversible eye damage. **DO NOT** get in eyes, on skin, or on clothing. **DO NOT breathe dust. DO NOT contaminate food or feed products.** Once a container has been attached to the hopper, it is recommended that it be used completely. Make sure the hoppers are emptied while still in the field. Refer to **STORAGE AND DISPOSAL** statement for further instructions.

Keep Out of Reach of Domestic Animals.
Not for Use or Storage In or Around Home

WORK SAFETY RULE

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SENSITIVITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposure should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal.

Before using this product, consult the National Pesticide Information Center for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of terbufos. The toll-free number for the National Pesticide Information Center is 1-800-858-7378.

If handled indoors, provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

DO NOT rub eyes or mouths with hands. If you feel sick in any way, STOP work and get help right away. See **FIRST AID** section.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are those made of waterproof material. If you want more options, follow the instructions for **Category A** on an EPA chemical-resistance category selection chart.

Loaders, applicators and other handlers using engineering controls must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

In addition to the above, loaders must also wear:

- Chemical-resistant gloves (such as those made of waterproof material)
- Chemical-resistant apron
- See engineering controls for additional requirements

IMPORTANT: All other handlers performing tasks such as spill clean up or equipment cleaning, for which engineering controls are not feasible must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Protective eyewear
- A respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter
- Chemical-resistant apron if exposed to concentrate

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash all PPE after each day's use.

DO NOT reuse contaminated clothing. Clothing and protection equipment heavily contaminated must be destroyed according to state and local regulations. **HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED**

ENGINEERING CONTROL STATEMENT

This product is formulated into an enclosed system (**LOCK'n LOAD®**) that meets the definition of a closed loading system as defined by the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(4)]. In addition to wearing the required PPE for loaders using engineering controls, loaders must be provided, must have immediately available, and must wear in case of a broken package, spill or equipment breakdown: coveralls, protective eyewear, chemical-resistant footwear plus socks, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter.

Applicators must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides for dermal protection. In addition, the applicator:

- Must wear the PPE required for handlers using engineering controls.
- Must either use an enclosed cab that also provides equivalent respiratory protection to an organic vapor-removing respirator or wear a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23G), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter.
- Must be provided with, must have immediately available for use, and must wear if they must exit the cab in the treated area: protective eyewear, coveralls, chemical-resistant gloves, chemical-resistant footwear plus socks, and if using an enclosed cab that provides respiratory protection respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge, or canister with any N, R, P or HE prefilter.
- Must take off any PPE that was worn in the treated area before reentering the cab.
- Must store all such PPE in a chemical-resistant container such as a plastic bag, to prevent contamination inside of the cab.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FAILURE TO FOLLOW PRECAUTIONS INCLUDING WEARING PROPER PERSONAL PROTECTIVE EQUIPMENT (PPE) MAY RESULT IN SERIOUS OR EVEN LIFE-THREATENING POISONING REQUIRING IMMEDIATE MEDICAL ATTENTION. THE ACTIVE INGREDIENT OF THIS GRANULAR FORMULATION CAN BE ABSORBED THROUGH THE SKIN TO CAUSE POISONING.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish and wildlife and has been known to cause fish kills. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wastewater or rinsate. See **DIRECTIONS FOR USE** for required buffer zones or setbacks. Birds and mammals may be killed if granules are not properly covered with soil in all areas of the treated field and in loading areas. Cover, incorporate or clean up product that is spilled during loading or that is visible on the soil surface in turn areas.

DO NOT apply in wet soil conditions that may prevent the equipment from covering pesticide granules. **DO NOT** apply to frequently flooded areas.

Under certain field conditions, terbufos and its degradates have a potential for runoff into surface water for several weeks following application. Keep out of lakes, ponds, and streams.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, whether directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. This label must be in the possession of the user at the time of pesticide application.

BEFORE USING, READ PRECAUTIONARY STATEMENTS AND ENVIRONMENTAL HAZARDS.

Loader and applicator requirements to minimize the potential for runoff to surface water:

- Maintain a 500 ft vegetative buffer between treated area and surface water on neighboring land.
- Maintain a 500 ft buffer between a standpipe drain outlet and surface water on neighboring land.
- Maintain a 66 ft setback between the treated area and entry points to surface water bodies on non-highly erodible soils.
- Maintain a 300 ft setback between the treated area and entry points to surface water bodies on highly erodible soils (HES).
- Maintain a 66 ft setback between treated area and standpipes on terraced fields as well as a 66 ft vegetative buffer between the tile outlet and entry points to surface water bodies.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of **48 hours**. The **48-hour** REI is increased to **72 hours** in outdoor areas where average rainfall is less than 25 inches a year.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

AGRICULTURAL USE REQUIREMENTS (CONTINUED)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves (such as those made of waterproof material)
- Chemical-resistant footwear plus socks
- Protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

All setbacks must be planted to a crop or seeded with grass or other suitable cover, and all vegetative buffers must be seeded with grass or other suitable cover.

DO NOT load, rinse, or wash equipment within 300 ft from surface water bodies.

DO NOT load, rinse, or wash equipment within 50 ft from wells unless conducted on an impervious surface.

To prevent channeling of surface water runoff, adjust the planter row-cleaners appropriately to prevent rows lower in height than adjacent soil.

Crops	Pests Controlled	Rate of Counter® 15G Systemic Insecticide Nematicide	Application	Remarks
FIELD CORN, FIELD CORN GROWN FOR SEED, POPCORN AND SWEET CORN At Planting	Corn rootworms ¹ Wireworms White grubs Seedcorn maggots Seedcorn beetles Corn flea beetles Maize billbugs Southern corn billbugs Thrips Symphylans Chinchbugs ² Nematodes: Lance Lesion Root Knot Spiral Stunt Sting Stubby root Dagger Suppression of: Cutworms ³ Lesser cornstalk borers	Banded or In-Furrow 6.0 ⁴ to 8 oz. per 1,000 ft. of row for any row spacing. DO NOT exceed 8.7 pounds of Counter 15G per acre.	Banded Place granules in a 7 inch band over the row, in front of the presswheel and incorporate evenly into the top 1 inch of the soil. In-Furrow Place granules directly in the seed furrow behind the planter shoe.	For use on conventional, conservation, or no-till corn. In situations where crop debris or winds can prevent proper placement of granules, in-furrow applications are recommended. In-furrow applications reduce the potential for insecticide runoff resulting from excessive rain. The presswheel should be adjusted so that the closed seed furrow is higher than the surrounding soil to reduce the potential for insecticide run-off resulting from excessive rain. ¹ Counter 15G controls species of corn rootworms including Southern, Northern, Western, and Mexican corn rootworms. ² For use in the states of Alabama, Florida, Georgia, Louisiana, Mississippi and Texas only for early season control of light to moderate infestations. ³ Under dry soil conditions or heavy infestations, it may be necessary to apply an insecticide rescue treatment with another registered insecticide after corn emergence to control surviving cutworm larvae. ⁴ The 6.0 oz. rate should only be used on first-year corn where heavy rootworm population pressures are anticipated or for control of other pests. If application is made at planting, do not make postemergence or cultivation time treatments of Counter 15G ALS-inhibiting herbicides* SHOULD NOT be used if Counter 15G has been applied to corn at the time of planting. See additional information for "at cultivation" applications.
Postemergence Incorporated (Same as crops above)	Maize billbugs Southern corn billbugs	Banded 8 oz. per 1,000 ft. of row for any row spacing. DO NOT exceed 8.7 pounds of Counter 15G per acre.	Apply in a 7 inch band over the row of seedling corn plants when billbugs or damage are observed, usually in the 1 to 6 leaf stage of growth. Use cultivators or other suitable implement to lightly incorporate granules into soil.	Only one at-planting, postemergence incorporated, or cultivation-time treatment of Counter 15G may be used. Consult your state experiment station, state extension service or certified pest management specialist for proper timing of application.

Crops	Pests Controlled	Rate of Counter® 15G Systemic Insecticide Nematicide	Application	Remarks
At Cultivation (Same as crops above)	Corn rootworms	8 oz. per 1,000 ft. of row for any row spacing. DO NOT exceed 8.7 pounds of Counter 15G per acre.	Apply granules to base of plants just ahead of cultivation shovels so as to cover granules with soil.	DO NOT graze or cut for forage within 30 days of treatment. DO NOT harvest sweet corn within 60 days after application. For Field Corn Only: (Excluding Field Corn Grown For Seed) Counter 15G may be applied 7 days after application of ALS-inhibiting herbicides*.
SUGAR BEETS At Planting	Sugar beet root maggots ¹ Sugar beet leafhopper ² Aphids ³ Wireworms ⁴ White Grubs ⁵ Crown borer	Banded or Modified In-Furrow 4-8 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 13.1 lbs. of Counter 15G per acre.	Modified In-Furrow Apply in-furrow at planting time 2-3 inches behind the seed drop zone after some soil has covered the seed.	At Planting DO NOT place granules in direct contact with the seed or crop injury may occur. Power incorporation may be used. DO NOT incorporate deeper than 2 inches. ¹ If especially heavy infestations are expected, use the 8 oz. rate. ² Modified in-furrow only at the 8 oz. rate. Only one application of Counter 15G per year may be made.
Suppression of: Cutworms		Banded 8 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 13.1 lbs. of Counter 15G per acre.	Banded Apply in 5 to 7 inch band over the row and lightly incorporate into the soil with drag chains or tines.	
Suppression of: Sugar beet cyst nematode Control of: Sugar beet leafhopper Aphids		Knifed-in 8 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 13.1 lbs. of Counter 15G per acre.	Knifed-in Drill granules 2 inches into the side of the seed and 2 to 4 inches below the seed.	
SUGAR BEETS Postemergence	Sugar beet root maggots ¹ Sugar beet leafhopper Aphid Sugar beet root aphid Crown borer ²	Banded 4-8 oz. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded Apply in 5 to 7 inch band over the row and lightly incorporate into the soil with cultivation shovels or other suitable implement.	Postemergence For sugar beet root maggot control apply at the time of first fly emergence. DO NOT harvest sugar beets or feed tops to livestock within 110 days after at planting banded, at planting modified in-furrow or postemergence banded applications. DO NOT harvest sugar beets or feed tops to livestock within 150 days of a knifed-in application ¹ Postemergence – apply Counter 15G at the first sign of crown borer infestation.
GRAIN SORGHUM At Bedding	Greenbugs Corn leaf aphids	Knifed-in 7 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 11.3 lbs. of Counter 15G per acre.	Knifed-in Drill granules 1-4 inches below the seed OR 1-4 inches below the seed and up to 5 inches to the side of the seed.	DO NOT place granules in direct contact with seed as crop injury may occur. Only one application of Counter 15G per year may be used.
GRAIN SORGHUM At Planting	Corn rootworms including: Southern corn rootworms Wireworms White Grubs Nematodes Greenbugs Corn leaf aphids Chinchbugs ³	Banded 7 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 11.3 lbs. of Counter 15G per acre.	Banded* Place granules in a 5 to 7 inch band directly behind the planter shoe in front of the press wheel and lightly incorporate with drag chains or tines.	*DO NOT use banded applications for aphid control in New Mexico, Arkansas, Florida, Georgia, Louisiana, Mississippi and Texas only for early season control of light to moderate infestations. DO NOT graze or harvest grain sorghum forage within 50 days after treatment. DO NOT harvest grain sorghum fodder or grain within 100 days after treatment. ³ For use in the states of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi and Texas only for early season control of light to moderate infestations.
	Greenbugs Corn leaf aphids	Knifed-in 7 oz. per 1,000 ft. of row for any row spacing (minimum 20 inch row spacing) or no more than 11.3 lbs. of Counter 15G per acre.		DO NOT place granules in direct contact with seed as crop injury may occur. Only one application of Counter 15G per year may be used.

*Such as **Accent**®, **Accent Gold**®, **Basis Gold**®, **Steadfast**®, **Beacon**®, **Exceed**®, **Hornet**™, **Permit**™, **Scorpion III**®, **Celebrity**®, **Celebrity Plus**, **Lightning**®, **Northstar**™, and **Spirit**™ herbicides.

Use Rates as Pounds per Acre*			
(based on various row spacings and application rates)			
Row Width (Inches)	Application Rates (oz/1,000 ft. of row)		
	4 oz.	6 oz.	8 oz.
40"	3.3	4.9	6.5
38"	3.4	5.2	6.9
36"	3.6	5.4	7.3
34"	3.9	5.6	7.7
32"	4.1	6.1	8.2
30"	4.4	6.5	8.7
28"	4.7	7.1	9.5
26"	5.0	7.6	10.2
24"	5.4	8.2	10.9
22"	5.9	8.9	11.9
20"	6.5	9.8	13.1

*See directions for specific use rates on each crop.

CALIBRATION INFORMATION			
FIRST READ THIS LABEL			
It is important the applicator equipment be properly set to deliver the labeled rate. This chart will help determine the desired rate of application. When label is 8 ounces per 1,000 feet of row, use these suggested starting gauge settings, regardless of row spacing:			
APPLICATOR	PLANTING SPEED		
	3 MPH	5 MPH	7 MPH
John Deere Max-Emerge 2 Models 7200 and 7300	16	23	35
John Deere Max-Emerge Models 7000 and 7100 Odd Notches 5-15-25-35	11	15	19
John Deere Max-Emerge Models 7000 and 7100 Even Notches 0-10-20-30	13	19	25
John Deere 71 Flexi-Planter Metal Hoppers	1/28	2/6.0	2/15
International Harvester	1/7.0	2/1.5	2/7.5
New International Harvester	1/9.0	2/6.0	3/2.0
Gandy	18	24	28
New Noble	8	12	16
Old Noble	11	17	23
Allis Chalmers 78 and 79 Series	All speeds – Gauge 7		
Allis Chalmers #385	10	14	18
Buffalo (Gauged)	9	15	20
Buffalo (Flute-Roll Feed)	All speeds – 5/8 Revolutions Hexagonal Nut		
White	Refer to Noble Settings		

NOTE: These settings should be used as starting points only. Continually check the amount of Counter[®] 15 G Systemic Insecticide Nematicide used against the known length of row acreage and make further adjustments accordingly. Also, check calibration occasionally to make sure equipment wear, changing moisture conditions, etc., have not caused a change in flow rate.

STORAGE AND DISPOSAL
DO NOT contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store pesticide products in a secure locked area where children, unauthorized persons, and animals cannot enter.
DO NOT store in the same area with food or feed. DO NOT store open containers.
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance. Open dumping is prohibited.
CONTAINER DISPOSAL: Return empty container to point of purchase.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

Counter, LOCK'nLOAD and SMARTBOX are registered trademarks of Amvac Chemical Corporation.

Celebrity and Lightning are registered trademarks of BASF Corporation.

Accent, Accent Gold, Basis Gold and Steadfast are registered trademarks of E.I. DuPont de Nemours & Co., Inc.

Beacon and Exceed are registered trademarks and Northstar and Spirit are trademarks of Syngenta Crop Protection, Inc.

Hornet is a trademark and Scorpion III is a registered trademark of Dow Agrosciences LLC.

Permit is a registered trademark of Nissan Chemical Industries, Ltd.

AMVAC Chemical Corporation
4100 E. Washington Boulevard
Los Angeles, CA 90023 U.S.A.
1-323-264-3910
www.amvac-chemical.com



4100 E. Washington Boulevard
Los Angeles, CA 90023, U.S.A.

1-323-264-3910 • www.amvac-chemical.com